arXiv:1112.2771v1 [q-bio.GN] 13 Dec 2011

Global transposable characteristics in the yeast complete DNA sequence

Zuo-Bing Wu*
State Key Laboratory of Nonlinear Mechanics,
Institute of Mechanics,
Chinese Academy of Sciences, Beijing 100190, China

December 14, 2011

Abstract

Global transposable characteristics in the complete DNA sequence of the Saccharomyces cevevisiae yeast is determined by using the metric representation and recurrence plot methods. In the form of the correlation distance of nucleotide strings, 16 chromosome sequences of the yeast, which are divided into 5 groups, display 4 kinds of the fundamental transposable characteristics: a short period increasing, a long quasi-period increasing, a long major value and hardly relevant.

Keywords Yeast, DNA sequences, Coherence structure, Metric representation, Recurrence plot

---

[1] Correspondence to: wuzb@lnm.imech.ac.cn

1 Introduction

The recent complete DNA sequences of many organisms are available to systematically search of genome structure. For the large amount of DNA sequences, developing methods for extracting meaningful information is a major challenge for bioinformatics. To understand the one-dimensional symbolic sequences composed of the four letters 'A', 'C', 'G' and 'T' (or 'U'), some statistical and geometrical methods were developed[1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. In special, chaos game representation (CGR)[12], which generates a two-dimensional square from a one-dimensional sequence, provides a technique to visualize the composition of DNA sequences. The characteristics of CGR images was described as genomic signature, and classification of species in the whole bacteria genome was analyzed by making an Euclidean metric between two CGR images[13]. Based on the genomic signature, the distance between two DNA sequences depending on the length of nucleotide strings was presented[14] and the horizontal transfers in prokaryotes and eukaryotes were detected and charaterized[15, 16]. Recently, a one-to-one metric representation of the DNA sequences[17], which was borrowed from the symbolic dynamics, makes an ordering of subsequences in a plane. Suppression of certain nucleotide strings in the DNA sequences leads to a self-similarity of pattern seen in the metric representation of DNA sequences. Self-similarity limits of genomic signatures were determined as an optimal string length for generating the genomic signatures[18]. Moreover, by using the metric representation method, the recurrence plot technique of DNA sequences was established and employed to analyze correlation structure of nucleotide strings[19].

As a eukaryotic organism, yeast is one of the premier industrial microorganisms, because of its essential role in brewing, baking, and fuel alcohol production. In addition, yeast has proven to be an excellent model organism for the study of a variety of biological problems involving the fields of genetics, molecular biology, cell biology and other disciplines within the biomedical and life sciences. In April 1996, the complete DNA sequence of the yeast (Saccharomyces cevevisiae) genome, consisting of 16 chromosomes with 12 million basepairs, had been released to provide a resource of genome information of a single organism. However, only 43.3% of all 6000 predicted genes in the Saccharomyces cerevisiae yeast were functionally characterized when the complete sequence of the yeast genome became available[20]. Moreover, it was found that DNA transposable elements have ability to move from place to place and make many copies within the genome via the transposition[21, 22]. Therefore, the yeast complete DNA sequence remain a topic to be studied respect to its genome architecture structure in the whole sequence.

In this paper, using the metric representation and recurrence plot methods, we analyze global transposable characteristics in the yeast complete DNA sequence, i.e., 16 chromosome sequences.

2 Metric representation and recurrence plot methods

For a given DNA sequence $s_1 s_2 \cdots s_i \cdots s_N$ ($s_i \in \{A, C, G, T\}$), a plane metric representation is generated by making the correspondence of symbol s_i to number μ_i or $\nu_i \in \{0, 1\}$ and calculating values (α, β) of all subsequences $\Sigma_k = s_1 s_2 \cdots s_k$ ($1 \leq k \leq N$) defined as follows

$$\begin{aligned} \alpha &= 2\sum_{j=1}^{k} \mu_{k-j+1} 3^{-j} + 3^{-k} = 2\sum_{i=1}^{k} \mu_i 3^{-(k-i+1)} + 3^{-k}, \\ \beta &= 2\sum_{j=1}^{k} \nu_{k-j+1} 3^{-j} + 3^{-k} = 2\sum_{i=1}^{k} \nu_i 3^{-(k-i+1)} + 3^{-k}, \end{aligned} \tag{1}$$

where μ_i is 0 if $s_i \in \{A, C\}$ or 1 if $s_i \in \{G, T\}$ and ν_i is 0 if $s_i \in \{A, T\}$ or 1 if $s_i \in \{C, G\}$. Thus, the one-dimensional symbolic sequence is partitioned into N subsequences Σ_k and mapped in the two-dimensional plane (α, β). Subsequences with the same ending l-nucleotide string, which are labeled by Σ^l, correspond to points in the zone encoded by the l-nucleotide string. Taking a subsequence $\Sigma_i \in \Sigma^l$, we calculate

$$\Theta(\epsilon_l - |\Sigma_i - \Sigma_j|) = \Theta(\epsilon_l - \sqrt{(\alpha_i - \alpha_j)^2 + (\beta_i - \beta_j)^2}), \tag{2}$$

where Θ is the Heaviside function $[\Theta(x) = 1$, if $x > 0$; $\Theta(x) = 0$, if $x \leq 0]$ and Σ_j is a subsequence $(j \geq l)$. When $\Theta(\epsilon_l - |\Sigma_i - \Sigma_j|) = 1$, i.e., $\Sigma_j \in \Sigma^l$, a point (i, j) is plotted in a plane. Thus, repeating the above process from the beginning of one-dimensional symbolic sequence and shifting forward, we obtain a recurrence plot of the DNA sequence.

For presenting correlation structure in the recurrence plot plane, a correlation intensity is defined at a given correlation distance d

$$\Xi(d) = \sum_{i=1}^{N-d} \Theta(\epsilon_l - |\Sigma_i - \Sigma_{i+d}|). \tag{3}$$

The quantity displays the transference of l-nucleotide strings in the DNA sequence. To further determine positions and lengths of the transposable elements, we analyze the recurrent plot plane. Since Σ_i and $\Sigma_j \in \Sigma^l$, the transposable element has the length l at least. From the recurrence plot plane, we calculate the maximal value of x to satisfy

$$\Theta(\epsilon_l - |\Sigma_{i+x} - \Sigma_{j+x}|) = 1, \quad x = 0, 1, 2, \cdots \tag{4}$$

i.e., Σ_{i+x} and $\Sigma_{j+x} \in \Sigma^l$. Thus, the transposable element with the correction distance $d = j - i + 1$ has the length $L = l + x$. The transposable element is placed at the position $(i - l + 1, i + x)$ and $(j - l + 1, j + x)$.

3 Global transposable characteristics in the yeast complete DNA sequence

The Saccharomyces cevevisiae yeast has 16 chromosomes, which are denoted as YEAST I to XVI. Using the metric representation and recurrence plot methods, we analyze correlation structures of the 16 DNA sequences. According to the characteristics of the correlation structures, we summarize the results as follows:

(1) The correlation distance has a short period increasing. The YEAST I, IX and XI have such characteristics. Let me take the YEAST I as an example to analyze. Fig.1 displays the correlation intensity at different correlation distance $d(\leq N-l)$ with $l=15$. A local region is magnified in the figure. It is clearly evident that there exist some equidistance parallel lines with a basic correlation distance $d_b=135$. Using Eq. (4), we determine positions and lengths of the transposable elements in Table I, where their lengths are limited in $L\geq 100$. Many nucleotide strings have correlation distance, which is the integral multiple of d_b. They mainly distribute in two local regions of the DNA sequence (25715-26845) and (204518-206554) or (11.2-11.7%) and (88.8-89.7%) expressed as percentages. The YEAST IX and XI have similar behaviors. The YEAST IX has the basic correlation distance $d_b=18$. Many nucleotide strings ($L\geq 50$) with the integral multiple of d_b mainly distribute in a local region of the DNA sequence (391337-393583) or (89.0-89.5%) expressed as percentages. The YEAST XI has the basic correlation distance $d_b=189$. Many nucleotide strings ($L\geq 50$) with the integral multiple of d_b mainly distribute in a local region of the DNA sequence (647101-647783) or (97.1-97.2%) expressed as percentages.

(2) The correlation distance has a long major value and a short period

increasing. The YEAST II, V, VII, VIII, X, XII, XIII, XIV, XV and XVI have such characteristics. Let me take the YEAST II as an example to analyze. Fig.2 displays the correlation intensity at different correlation distance $d(\leq N-l)$ with $l=15$. The maximal correlation intensity appears at the correlation distance $d_m = 38534$. A local region is magnified in the figure. It is clearly evident that there exist some equidistance parallel lines with a basic correlation distance $d_b = 36$. In Table II, positions and lengths $(L \geq 50)$ of the transposable elements are given. The maximal transposable elements mainly distribute in two local regions of the DNA sequence (221249-224565, 259783-263097) or (27.2-27.6%, 31.9- 32.4%) expressed as percentage. Near the positions, there also exist some transposable elements with approximate values for d_b. Moreover, many nucleotide strings have correlation distance, which is the integral multiple of d_b. They mainly distribute in a local region of the DNA sequence (391337-393583) or (89.0-89.5%) expressed as percentages. In the other 9 DNA sequences, the YEAST V, X, XII, XIII, XIV, XV and XVI have the same basic correlation distance $d_b = 36$ and similar behaviors with different major correlation distance d_m =49099, 5584, 9137, 12167, 5566, 447110 and 45988, respectively. The YEAST VII and VIII have different basic correlation distance $d_b = 12$ and 135, and similar behaviors with the major correlation distance $d_m = 255548$ and 1998, respectively.

(3) The correlation distance has a long quasi-period increasing. The YEAST III has such characteristics. Fig. 3 displays the coherence intensity at different correlation distance $d(\leq N-l)$ with $l=15$. The correlation intensity has the maximal value at the correlation distance $d_{m1} = 185903$ and two vice-maximal values at the correlation distance $d_{m2} = 93625$ and $d_{m3} = 279528$. Since $d_{m2} \approx d_{m1}/2 \approx d_{m3}/3$, the coherence distance has a quasi-period increasing. A local region is magnified in the figure. These

does not exist any clear short period increasing of the correlation distance. Using Eq. (4), we determine positions and lengths ($L \geq 50$) of the transposable elements in Table III. The maximal and vice-maximal transposable elements mainly distribute in local regions of the DNA sequence (11499-13810, 197402-199713), (198171-199796, 291794-293316) and (12268-12932, 291794-292460) or (3.6-4.4%, 62.6-63.6%), (62.8-63.4%, 92.5-93.0%) and (3.9-4.1%, 92.5-92.7%) expressed as percentage.

(4) The correlation distance has a long major value and a long quasi-period and two short period increasing. The YEAST IV has such characteristics. Fig. 4 displays the coherence intensity at different correlation distance $d(\leq N-l)$ with $l=15$. The maximal coherence intensity appears at the correlation distance $d_m = 3885$. There also exist three vice-maximal values at the correlation distance $d_{m1} = 232800$, $d_{m2} = 109349$ and $d_{m3} = 341221$, which forms a long quasi-period increasing of the correlation distance, i.e., $d_{m2} \approx d_{m1}/2 \approx d_{m3}/3$. A local region is magnified in the figure. It is clearly evident that there exist two short period increasing with $d_{b1} = 84$ and $d_{b2} = 192$ in the correlation distance. In Table IV, positions and lengths ($L \geq 100$) of the transposable elements are determined by using Eq. (4). All correlation distance with the long major value and the long quasi-period and two short period increasing are denoted. The transposable elements with d_m, d_{m1}, d_{m2}, d_{m3}, d_{b1} and d_{b2} mainly distribute in local regions of the DNA sequence (527570-538236), (871858-876927, 981207-986276), (645646-651457, 878346-884257), (646379-651032, 987600-992253), (1307733-1308591) and (758135-759495) or (34.4-35.1%), (56.9-57.2%, 64.0-64.4%), (42.1-42.5%, 57.3-57.7%), (42.2-42.5%, 64.4-64.8%), (85.36-85.41%) and (49.5-49.6%) expressed as percentages.

(5) The DNA sequence is hardly relevant. The YEAST VI has such

characteristics. Fig. 5 displays the coherence intensity at different correlation distance $d(\leq N-l)$ with $l = 15$. The maximal coherence intensity appears at the correlation distance $d_m = 5627$. A local region is magnified in the figure. The sequence has not a short period increasing of the coherence distance. In Table V, positions and lengths ($L \geq 50$) of the transposable elements are given. Only one nucleotide string with the length 337 has the correlation distance d_m. The YEAST VI is almost never relevant, so the YEAST VI approaches a random sequence.

4 Conclusion

Global transposable characteristics in the yeast complete DNA sequence is determined by using the metric representation and recurrence plot methods. Positions and lengths of all transposable nucleotide strings in the 16 chromosome DNA sequences of the yeast are determined. In the form of the correlation distance of nucleotide strings, the fundamental transposable characteristics displays a short period increasing, a long quasi-period increasing, a long major value and hardly relevant. The 16 chromosome sequences are divided into 5 groups, which have one or several of the 4 kinds of the fundamental transposable characteristics.

Acknowledgments We thank the IMECH and ICTS research computing facilities for assisting us in the computation.

References

[1] R. Durbin, S.R. Eddy, A. Krogh, and G. Mitchison. *Probabilistic models of proteins and nucleic acids* (1999), Cambridgr University Press.

[2] B.-L. Hao, H. C. Lee, and S.-Y. Zhang. Fractals related to long DNA sequences and bacterial complete genomes. *Chaos, Solitons and Fractals* **11** (2000) 825.

[3] D. Qi, A. J. Cuticchia, Compositional symmetries in complete genomes. *Bioinformatics* **17** (2001) 557.

[4] W. Li, P. Bernaola-Galván, F. Haghighi, I. Grosse. Applications of recursive segmentation to the analysis of DNA sequences. *Computers & Chemistry* **26** (2002) 491.

[5] D. Robelin, H. Richard, B. Prum, SIC: a tool to detect short inverted segments in a biological sequence, *Nucleic Acids Research* **31** (2003) 3669.

[6] S. Garte, Fractal properties of the human genome. *Journal of Theoretical Biology* **230** (2004) 251.

[7] P. W. Messer, P. F. Arndt, M. Lässig (2005), Solvable sequence evolution models and genomic correlations, *Phys. Rev. Lett.* **94** (2005) 138103.

[8] W. Li, P. Miramontes, Large-scale oscillation of structure-related DNA sequence features in human chromosome 21, *Physical Review E* **74** (2006) 021912.

[9] A. Provata, Th. Oikonomou, Power law exponents characterizing human DNA, *Physical Review E* **75** (2007) 056102.

[10] H. Gonzalez-Diaz, Y. Gonzalez-Diaz, L. Santana, et al. Proteomics, networks and connectivity indices, *Proteomics* **8** (2008) 750.

[11] J. S. Almeida and S. Vinga, Biological sequences as pictures - a generic two dimensional solution for iterated maps, *BMC Binoinformatics* **10** (2009) 100.

[12] H. J. Jeffrey, Chaos game representation of gene structure. *Nucleic Acids Res.* **18** (1990) 2163.

[13] P. J. Deschavanne, A. Giron, J. Vilain, G. Fagot, and B. Fertil, Genomic signature: characterization and classification of species assessed by chaos game representation of sequences. *Mol. Biol. Evol.* **16** (1999) 1391.

[14] Y. Wang, K. Hill, S. Singh and L. Kari, The spectrum of genomic signatures: from dinucleotides to chaos game representation. *Gene* **346** (2005) 173.

[15] C. Dufraigne, B. Feitil, S. Lespinats, A. Giron and P. Deschavanne, Detection and characterization of horizontal transfers in prokaryotes using genomic signature. *Nucleic Acids Res.* **33** (2005) e6.

[16] L. V. Mallet, J. Becq and P. Deschavanne, Whole genome evaluation of horizontal transfers in the pathogenic fungus Aspergillus fumigatus. *BMC Genomics* **11** (2010) 171.

[17] Z.-B. Wu, Metric representation of DNA sequences. *Electrophoresis* **21** (2000) 2321.

[18] Z.-B. Wu, Self-similarity limits of genomic signatures. *Fractals* **11** (2003) 19.

[19] Z.-B. Wu, Recurrence plot analysis of DNA sequences. *Phys. Lett. A* **232** (2004) 250.

[20] H. W. Mewes, K. Albermann, M. Bähr et al. Overview of the yeast genome, *Nature* **387** (1997) 7.

[21] H. Ochman, J. G. Lawrence and E. A. Groisman, Lateral gene transfer and the nature of bacterial innovation, *Nature* **405** (2000) 299.

[22] J. L. Bennetzen, Transposable element contributions to plane gene and genome evolution, *Plant Mol. Biol.* **42** (2000) 251.

Table I. Transference of nucleotide strings with lengths $L(\geq 100)$ for the YEAST I with 230209 bases.

No.	String	Position 1	Position 2	L	d	Note
1	$t^2a\cdots act$	11745-11969	24177-24401	225	12432	
2	$ctg\cdots a^2t$	12258-12396	24711-24849	139	12453	
3	$g^2a\cdots g^2a$	12988-13171	25153-25336	184	12165	
4	$c^2g\cdots cgt$	25715-25851	26255-26391	137	540	$4d_b$
5	$at^2\cdots ac^2$	25739-25851	26414-26526	113	675	$5d_b$
6	$gta\cdots ac^2$	25751-25851	26561-26661	101	810	$6d_b$
7	$gta\cdots ac^2$	25751-25851	26696-26796	101	945	$7d_b$
8	$t^2g\cdots g^2t$	25853-25968	26393-26508	116	540	$4d_b$
9	$atg\cdots gtg$	25925-26035	26060-26170	111	135	d_b
10	$atg\cdots agt$	25925-26058	26195-26328	134	270	$2d_b$
11	$agt\cdots gtg$	26050-26170	26185-26305	121	135	d_b
12	$at^2\cdots gac$	26279-26406	26414-26541	128	135	d_b
13	$gta\cdots gac$	26291-26406	26561-26676	116	270	$2d_b$
14	$gta\cdots gac$	26291-26406	26696-26811	116	405	$3d_b$
15	$gta\cdots gtg$	26426-26710	26561-26845	285	135	d_b
16	$tga\cdots aca$	160239-160575	165827-166163	337	5588	
17	$cac\cdots tac$	204518-204802	204653-204937	285	135	d_b
18	$g^2t\cdots tac$	204567-204667	205512-205612	101	945	$7d_b$
19	$g^2t\cdots tac$	204702-204802	205512-205612	101	810	$6d_b$
20	$g^2t\cdots a^2t$	204837-204949	205512-205624	113	675	$5d_b$
21	$ac^2\cdots c^2a$	204855-204969	205395-205509	115	540	$4d_b$
22	$ctc\cdots cat$	205042-205168	205312-205438	127	270	$2d_b$
23	$cac\cdots act$	205058-205178	205193-205313	121	135	d_b
24	$cac\cdots cat$	205193-205303	205328-205438	111	135	d_b
25	$atg\cdots t^2c$	205758-205879	206433-206554	122	675	$5d_b$

Table II. Transference of nucleotide strings with lengths $L(\geq 50)$ for the YEAST II with 813142 bases. Due to the limited spacing, 22 nucleotide strings in the total number 57 are not presented.

No.	String	Position 1	Position 2	L	d	Note
1	$tag \cdots agt$	1584-1650	1692-1758	67	108	$3d_b$
2	$tag \cdots gct$	1620-1674	2016-2070	55	396	$11d_b$
3	$tag \cdots gct$	1620-1674	2268-2322	55	648	$18d_b$
4	$tgc \cdots tg^2$	1815-1870	1851-1906	56	36	d_b
5	$gct \cdots gta$	1860-1936	1932-2008	77	72	$2d_b$
6	$tgc \cdots gta$	1887-1936	2355-2404	50	468	$13d_b$
7	$tgc \cdots gtg$	1959-2018	2355-2414	60	396	$11d_b$
8	$gca \cdots agt$	2012-2082	2264-2334	71	252	$7d_b$
9	$tg^2 \cdots agt$	2022-2190	2094-2262	169	72	$2d_b$
10	$cag \cdots gta$	2037-2104	2325-2392	68	288	$8d_b$
11	$tg^2 \cdots agt$	2094-2154	2274-2334	61	180	$5d_b$
12	$cag \cdots gta$	2109-2176	2325-2392	68	216	$6d_b$
13	$tg^2 \cdots agt$	2166-2226	2274-2334	61	108	$3d_b$
14	$cag \cdots gta$	2181-2248	2325-2392	68	144	$4d_b$
33	$a^2t \cdots c^2t$	221182-221231	259718-259767	50	38536	$\approx d_m$
35	$gta \cdots tct$	221249-221308	259783-259842	60	38534	d_m
37	$gat \cdots tca$	222669-222827	261203-261361	159	38534	d_m
38	$ta^2 \cdots tgc$	222829-223096	261363-261630	268	38534	d_m
39	$cat \cdots gct$	223098-223609	261632-262143	512	38534	d_m
40	$a^2c \cdots g^2a$	223611-223966	262145-262500	356	38534	d_m
41	$tc^2 \cdots c^2g$	223968-224563	262502-263097	596	38534	d_m
42	$ga^2 \cdots tac$	224736-225308	263273-263845	573	38537	$\approx d_m$
43	$ata \cdots aca$	225310-225493	263847-264030	184	38537	$\approx d_m$
44	$ac^2 \cdots a^2c$	225841-226231	264378-264768	391	38537	$\approx d_m$
45	$acg \cdots t^3$	226233-226797	264770-265334	565	38537	$\approx d_m$

Table III. Transference of nucleotide strings with lengths $k(\geq 50)$ for the YEAST III with 315341 bases.

No.	String	Position 1	Position 2	L	d	Note
1	$c^3 \cdots ac^2$	90-141	233-284	52	143	
2	$a^2t \cdots ca^2$	1190-1253	4083-4146	64	2893	
3	$t^2g \cdots gta$	11499-11764	197402-197667	266	185903	d_{m1}
4	$tag \cdots cta$	11766-11908	197669-197811	143	185903	d_{m1}
5	$ta^2 \cdots gac$	11910-12185	197813-198088	276	185903	d_{m1}
6	$at^2 \cdots gtg$	12187-13810	198090-199713	1624	185903	d_{m1}
7	$tac \cdots tat$	12268-12340	291794-291866	73	279526	$\approx d_{m3}$
8	$ata \cdots at^2$	12325-12932	291853-292460	608	279528	d_{m3}
9	$a^2t \cdots gtg$	13691-13810	293114-293233	120	279423	
10	$cg^2 \cdots ca^2$	13812-14006	199715-199909	195	185903	d_{m1}
11	$cg^2 \cdots t^2c$	13812-13893	293235-293316	82	279423	
12	$tgt \cdots a^2c$	83677-83802	84419-84544	126	742	
13	$tgt \cdots a^2c$	83677-83802	90049-90174	126	6372	
14	$cta \cdots a^2c$	83724-83802	83951-84029	79	227	
15	$ca^2 \cdots tg^2$	83804-83902	84031-84129	99	227	
16	$ca^2 \cdots tg^2$	83804-83902	84546-84644	99	742	
17	$ca^2 \cdots tg^2$	83804-83902	90176-90274	99	6372	
18	$cta \cdots tct$	83951-84230	84466-84745	280	515	
19	$cta \cdots tat$	83951-84189	90096-90334	239	6145	
20	$tgt \cdots tat$	84419-84704	90049-90334	286	5630	
21	$cac \cdots t^3$	123942-124058	142661-142777	117	18719	
22	$tac \cdots tat$	198171-198243	291794-291866	73	93623	$\approx d_{m2}$
23	$ata \cdots at^2$	198228-198835	291853-292460	608	93625	d_{m2}
24	$a^2t \cdots t^2c$	199594-199796	293114-293316	203	93520	$\approx d_{m2}$
25	$g^2a \cdots g^3$	267365-267574	267692-267901	210	327	

Table IV. Transference of nucleotide strings with lengths $L(\geq 100)$ for the YEAST IV with 1531977 bases. Due to the limited spacing, 112 nucleotide strings without any notes in the total number 176 are not presented.

No.	String	Position 1	Position 2	L	d	Note
28	$gct\cdots a^3$	527570-527835	531455-531720	266	3885	d_m
29	$gct\cdots a^2t$	527570-527781	535340-535551	212	7770	
30	$gt^2\cdots c^2a$	527891-528066	531776-531951	176	3885	d_m
31	$gt^2\cdots c^2a$	527891-528066	535661-535836	176	7770	
32	$ag^2\cdots a^2t$	528094-531666	531979-535551	3573	3885	d_m
35	$ct^2\cdots t^2g$	531668-532364	535553-536249	697	3885	d_m
36	$gca\cdots c^2a$	532366-534351	536251-538236	1986	3885	d_m
40	$ata\cdots agt$	645546-646118	878346-878918	573	232800	d_{m2}
41	$ag^2\cdots aca$	645635-645873	992440-992678	239	346805	
42	$agc\cdots atc$	646120-646336	878920-879136	217	232800	d_{m2}
43	$ct^2\cdots gta$	646338-646726	879138-879526	389	232800	d_{m2}
44	$t^3\cdots atg$	646379-646564	987600-987785	186	341221	d_{m3}
45	$t^2c\cdots cat$	646787-647179	879587-879979	393	232800	d_{m2}
46	$tat\cdots t^2c$	646964-647288	988185-988509	325	341221	d_{m3}
47	$t^3\cdots ta^2$	647310-647470	880110-880270	161	232800	d_{m2}
48	$ta^2\cdots tga$	647468-647693	988689-988914	226	341221	d_{m3}
49	$gt^2\cdots cgt$	647472-648042	880272-880842	571	232800	d_{m2}
50	$gag\cdots aga$	647695-649519	988916-990740	1825	341221	d_{m3}
51	$gat\cdots cg^2$	648165-648265	880965-881065	101	232800	d_{m2}
52	$tc^2\cdots atg$	648486-649715	881286-882515	1230	232800	d_{m2}
53	$cgt\cdots atg$	649521-649715	990742-990936	195	341221	d_{m3}
54	$ctg\cdots t^3$	649853-650096	991074-991317	244	341221	d_{m3}
55	$tag\cdots ctc$	649946-650073	882746-882873	128	232800	d_{m2}
56	$gat\cdots aca$	650075-651457	882875-884257	1383	232800	d_{m2}
57	$ctg\cdots ct^2$	650098-651032	991319-992253	935	341221	d_{m3}

60	$ag^2\cdots aca$	651219-651457	992440-992678	239	341221	d_{m3}
63	$ac^2\cdots c^2a$	757478-757581	757670-757773	104	192	d_{b2}
68	$cta\cdots act$	758135-758259	758519-758643	125	384	$2d_{b2}$
69	$cta\cdots act$	758135-758259	758711-758835	125	576	$3d_{b2}$
70	$cta\cdots act$	758135-758259	759479-759603	125	1344	$7d_{b2}$
71	$gca\cdots g^2a$	758219-758343	758411-758535	125	192	d_{b2}
72	$gca\cdots a^2t$	758219-758347	759179-759307	129	960	$5d_{b2}$
73	$gca\cdots g^2a$	758219-758343	759371-759495	125	1152	$6d_{b2}$
74	$ctg\cdots act$	758349-758451	758925-759027	103	576	$3d_{b2}$
75	$ctg\cdots g^2a$	758349-758535	759117-759303	187	768	$4d_{b2}$
76	$ctg\cdots cat$	758349-758683	759309-759643	335	960	$5d_{b2}$
77	$cta\cdots cat$	758495-758683	758687-758875	189	192	d_{b2}
78	$cta\cdots aga$	758687-758901	759455-759669	215	768	$4d_{b2}$
79	$gca\cdots act$	758795-759027	758987-759219	233	192	d_{b2}
80	$cta\cdots act$	758903-759027	759287-759411	125	384	$2d_{b2}$
81	$gca\cdots aga$	758987-759093	759563-759669	107	576	$3d_{b2}$
82	$cta\cdots g^2a$	759095-759303	759287-759495	209	192	d_{b2}
99	$tgt\cdots a^3$	871858-872030	981207-981379	173	109349	d_{m1}
103	$tgc\cdots ca^2$	872202-872307	981551-981656	106	109349	d_{m1}
104	$a^2g\cdots cag$	872309-872592	981658-981941	284	109349	d_{m1}
105	$at^2\cdots tat$	872737-872871	982086-982220	135	109349	d_{m1}
106	$ag^2\cdots ca^2$	873022-873286	982371-982635	265	109349	d_{m1}
107	$tga\cdots cat$	873378-874087	982727-983436	710	109349	d_{m1}
108	$tca\cdots g^2t$	874089-874236	983438-983585	148	109349	d_{m1}
109	$tac\cdots tgc$	874238-874593	983587-983942	356	109349	d_{m1}
110	$aga\cdots atc$	874595-874764	983944-984113	170	109349	d_{m1}
111	$t^3\cdots ca^2$	874853-875247	984202-984596	395	109349	d_{m1}
112	$gat\cdots a^2c$	875249-875604	984598-984953	356	109349	d_{m1}

113	$ca^2 \cdots tga$	875637-876474	984986-985823	838	109349	d_{m1}
114	$agc \cdots tga$	876491-876927	985840-986276	437	109349	d_{m1}
117	$g^2a \cdots aga$	877085-877385	986434-986734	301	109349	d_{m1}
118	$ga^2 \cdots a^3$	877387-877657	986736-987006	271	109349	d_{m1}
169	$cgt \cdots ac^2$	1307733-1307835	1308321-1308423	103	588	$7d_{b1}$
170	$ac^2 \cdots g^2c$	1307749-1307874	1308505-1308630	126	756	$9d_{b1}$
171	$gt^2 \cdots atc$	1307753-1307871	1307921-1308039	119	168	$2d_{b1}$
172	$gt^2 \cdots atc$	1307921-1308039	1308509-1308627	119	588	$7d_{b1}$
173	$gt^2 \cdots atc$	1308089-1308249	1308341-1308501	161	252	$3d_{b1}$
174	$ac^2 \cdots atc$	1308169-1308333	1308253-1308417	165	84	d_{b1}
175	$ac^2 \cdots ac^2$	1308337-1308507	1308421-1308591	171	84	d_{b1}

Table V. Transference of nucleotide strings with lengths $L(\geq 50)$ for the YEAST VI with 270148 bases.

No.	String	Position 1	Position 2	L	d	Note
1	$tat \cdots aca$	137905-138238	143532-143865	334	5627	d_m
2	$ga^2 \cdots t^3$	178016-178086	178157-178227	71	141	
3	$ca^2 \cdots gtc$	178088-178157	178229-178298	70	141	
4	$tgt \cdots gtg$	210332-210391	210334-210393	60	2	

Figure caption

Fig. 1. A plot of correlation intensity $\Xi(d)$ versus correlation distance d for the YEAST I.

Fig. 2. A plot of correlation intensity $\Xi(d)$ versus correlation distance d for the YEAST II.

Fig. 3. A plot of correlation intensity $\Xi(d)$ versus correlation distance d for the YEAST III.

Fig. 4. A plot of correlation intensity $\Xi(d)$ versus correlation distance d for the YEAST IV.

Fig. 5. A plot of correlation intensity $\Xi(d)$ versus correlation distance d for the YEAST VI.